

The efficacy of intrauterine devices for emergency contraception: a systematic review of 35 years of experience

Kelly Cleland^{1,*}, Haoping Zhu², Norman Goldstuck³, Linan Cheng⁴,
and James Trussell^{1,5}

¹Office of Population Research, Princeton University, 218 Wallace Hall, Princeton, NJ 08544, USA ²Minhang Central Hospital, Shanghai Jiaotong University, Shanghai 201100, People's Republic of China ³Reproduction Research South Africa, St John's Rd, Sea Point, Cape Town 8005, South Africa ⁴Shanghai Institute of Planned Parenthood Research, Xie Tu Rd, Shanghai 200032, People's Republic of China ⁵The Hull York Medical School, University of Hull, Hull HU6 7RX, UK

*Correspondence address: Tel: +1-609-258-1395; Fax: +1-609-258-1039; E-mail: kcleland@princeton.edu

Submitted on February 1, 2012; resubmitted on March 19, 2012; accepted on March 23, 2012

BACKGROUND: Intrauterine devices (IUDs) have been studied for use for emergency contraception for at least 35 years. IUDs are safe and highly effective for emergency contraception and regular contraception, and are extremely cost-effective as an ongoing method. The objective of this study was to evaluate the existing data to estimate the efficacy of IUDs for emergency contraception.

METHODS: The reference list for this study was generated from hand searching the reference lists of relevant articles and our own article archives, and electronic searches of several databases: Medline, Global Health, Clinicaltrials.gov, Popline, Wanfang Data (Chinese) and Weipu Data (Chinese). We included studies published in English or Chinese, with a defined population of women who presented for emergency contraception and were provided with an IUD, and in which the number of pregnancies was ascertained and loss to follow-up was clearly defined. Data from each article were abstracted independently by two reviewers.

RESULTS: The 42 studies (of 274 retrieved) that met our inclusion criteria were conducted in six countries between 1979 and 2011 and included eight different types of IUD and 7034 women. The maximum timeframe from intercourse to insertion of the IUD ranged from 2 days to 10 or more days; the majority of insertions (74% of studies) occurred within 5 days of intercourse. The pregnancy rate (excluding one outlier study) was 0.09%.

CONCLUSIONS: IUDs are a highly effective method of contraception after unprotected intercourse. Because they are safe for the majority of women, highly effective and cost-effective when left in place as ongoing contraception, whenever clinically feasible IUDs should be included in the range of emergency contraception options offered to patients presenting after unprotected intercourse. This review is limited by the fact that the original studies did not provide sufficient data on the delay between intercourse and insertion of the IUD, parity, cycle day of intercourse or IUD type to allow analysis by any of these variables.

Key words: emergency contraception / intrauterine device / unintended pregnancy / unprotected intercourse / contraception

Introduction

Unintended pregnancy is a significant problem worldwide. It is estimated that globally at least 36% of pregnancies are unintended (Singh *et al.*, 2009), and in the USA nearly half of pregnancies are unintended (Finer and Zolna, 2011). Emergency contraception offers women an important strategy to prevent pregnancy after intercourse in cases of contraceptive accidents or non-use, or in situations of sexual violence. There are two forms of emergency contraception available today: pills and intrauterine devices (IUDs). The most

common medication option is 1.5 mg levonorgestrel, sold in one-pill or two-pill formulations. A newer formulation is 30 mg ulipristal acetate, marketed in the USA as ella[®] and in much of Europe as ellaOne[®]. In a few places, such as China, Vietnam and Russia, mifepristone in small doses is available for emergency contraception.

Non-hormonal IUDs (primarily copper-bearing) have been used for emergency contraception for at least 35 years (Lippes *et al.*, 1976). (The levonorgestrel intrauterine system, sold in the USA and Europe under the brand name Mirena[®], has not been studied for use for emergency contraception.) Negative experiences with the Dalkon

115 Shield, an IUD available in the 1970s in the USA, led to years of
concern about the safety of IUDs and very low levels of IUD use.
However, the design of modern IUDs available today is vastly
120 improved, and guidelines from major medical organizations, such as
the Centers for Disease Control and Prevention, the World Health
Organization, the UK Faculty of Sexual and Reproductive Healthcare
and American College of Obstetricians and Gynecologists, note that
IUDs are a safe choice for the majority of patients, including young
and nulliparous women (World Health Organization, 2009; ACOG
Practice Bulletin, 2010; Centers for Disease Control and Prevention,
125 2010; Faculty of Sexual & Reproductive Healthcare, Clinical Effective-
ness Unit, 2011).

One of the major advantages of copper IUDs is that following use
for emergency contraception, they can then be left in place to provide
at least 10 years of highly effective ongoing contraception. [In the USA
130 ParaGard[®] is labeled for 10 years of use but there is evidence of ef-
ficacy with longer use (Dean and Schwarz, 2011).] IUDs have been
shown to be among the most cost-effective methods of contraception
(Trussell et al., 2009); the fact that this is a 'forgettable' method that
does not require action on the part of the user means that there is
135 virtually no scope for user error.

IUDs are experiencing a moderate comeback after years of very
low uptake in the USA (Hubacher et al., 2011). In 2008 (the last
year for which data are available), 4.9% of American women at risk
of pregnancy reported using an IUD (Mosher and Jones, 2010). This
140 is a marked increase from the 0.7% of women at risk of pregnancy
choosing IUDs in 1995 (Mosher and Jones, 2010) but is still lower
than the use in Europe, where 10% of British women (data from
2009) (Lader, 2009) and 24% of French women (data from 2005)
(Moreau et al., 2008) at risk of pregnancy use IUDs. No comparable
145 statistic is available for IUD use among all women at risk of pregnancy
in China, but the Chinese National Population and Family Planning
Commission reported that 53% of married women using contracep-
tion used IUDs in 2009 (National Population and Family Planning
Commission, Population and Development Research Center of
150 China, 2010). IUD use is higher in China than in the world overall; a
2005 report noted that 43% of Chinese women using contraception
used IUDs, compared with 13% in the rest of the world (Salem,
2006). Guidelines for the use of IUDs for emergency contraception
typically recommend inserting the IUD within 5 days of unprotected
155 intercourse (ACOG Practice Bulletin, 2010), although the Centers
for Disease Control, the World Health Organization and the UK
Faculty of Sexual and Reproductive Healthcare specify that an IUD
can be used beyond 5 days, as long as the time of ovulation can be
reasonably determined and the insertion occurs no more than 5
160 days after ovulation (World Health Organization, 2009; Centers for
Disease Control and Prevention, 2010; Faculty of Sexual & Reproduct-
ive Healthcare, Clinical Effectiveness Unit, 2011). It should be noted
that the guidelines around the time of insertion are not related to ef-
ficacy or safety but to ensure that the IUD is inserted before the im-
165 plantation of an embryo (thus ensuring its function as a contraceptive,
rather than an early abortifacient).

Materials and Methods

170 This study is a systematic review designed to provide a current estimate of
the efficacy of IUDs used for emergency contraception, based on all of the

available published data. The reference list for this study was generated
from searching our own archives, references lists of relevant articles and
queries of several databases using the following search terms:

MEDLINE: 'Contraception, Postcoital' [Mesh] AND 'Intrauterine
175 Devices' [Mesh],

Clinicaltrials.gov: 'intrauterine device' AND 'emergency contraception',
Popline: 'IUD' & 'Emergency Contraception',

Global Health: 'intrauterine device' AND 'emergency contracept*' OR
'postcoit* contracept*', 180

Wanfang Data (Chinese): (emergency contraception, intrauterine
device),

Weipu Data (Chinese): (emergency contraception, intrauterine device).

This review includes any peer-reviewed study published by August 2011
185 in English or Chinese, with a defined population of women who presented
for emergency contraception and were provided with an IUD, and in
which the number of pregnancies was ascertained and loss to follow-up
was clearly defined. Chinese data were included because of the tremen-
dous amount of contraceptive research taking place in that country.

Once relevant articles were identified, data from each study were
190 abstracted independently by two reviewers using a common data entry
form that captured the language of publication, country in which data
were collected, type(s) of IUD used, maximum time from intercourse to
IUD insertion, initial study enrollment, efficacy-evaluable population, loss
to follow-up and number of treatment failures (pregnancies).

We computed Blyth–Still–Casella exact 95% binomial confidence
195 intervals (CI) for proportions and used either Fisher's Exact Test (for
 2×2) or Fisher–Freeman–Halton (for $R \times 2$) to test for homogeneity.
All calculations were performed in StatXact in Cytel Studio 8 (Cytel
Inc., Cambridge, MA, USA). 200

Results

Our search found 274 articles, and we assessed 48 of these in depth
205 to determine their eligibility for inclusion. Five studies were excluded
from our analysis because they did not provide sufficient detail
about the loss to follow-up (Bromwich and Parsons, 1982; Hutchin-
son, 1983; Wright and Thompson, 1986; Friedman and Rowley,
1987; Zhang and Huang, 2005). In addition, one study was published
210 both in English and Chinese, and we included the English version in this
analysis (Fan and Zhou, 2001; Zhou and Xiao, 2001). This study
included a subject who presented 95 h after unprotected intercourse
and was believed to be pregnant as a result of an act of intercourse 16
215 days prior to insertion of the IUD; this individual was excluded from
the analysis. One study was designed to compare the insertion toler-
ability of the GyneFix IUD versus the Gyne-T3805 IUD for emergency
contraception but it did report that no pregnancies occurred in the
trial, and so we included it in our final dataset (D'Souza et al.,
2003). Forty-two studies were included in the final review.

The 42 studies that fit our eligibility criteria and were included in the
220 review ranged in the year of publication from 1979 to 2011 (Table 1).
Of these, 28 were published in Chinese and 14 in English. The English
literature included data collected in China, Egypt, Italy, the Nether-
lands, USA and the UK. Nearly all of the IUDs were copper-bearing,
225 although a small number of plastic IUDs (the Lippes Loop series) were
used in two of the earlier studies (Black et al., 1980; Guillebaud et al.,
1983). The majority of studies in our review (31 studies, 74%) fol-
lowed the current standard protocol of inserting the IUD within 5

Table 1 Studies included in review of the efficacy of IUDs for emergency contraception over 35 years, in order of publication.

Study	Country	IUD type	Max. time to insertion (days)	Enrolled (n)	Efficacy population (n)	Pregnancies (n)	Pregnancy rate (%) (95% CI) ^a
Gottardi <i>et al.</i> (1979) ^b	Italy	Cu T-200	2	60	60	0	0 (0–6.0)
Lippes <i>et al.</i> (1979) ^b	USA	Cu T-200, Cu 7	7	299	299	0	0 (0–1.2)
Tyrer (1980) ^b	USA	Cu 7	5	80	80	0	0 (0–4.5)
Black <i>et al.</i> (1980) ^b	UK	Cu 7, Lippes Loop D	10	191	176	0	0 (0–2.1)
Van Santen and Haspels (1981) ^b	Netherlands	MLCu-250, Cu 7, Cu T-200	5	55	55	0	0 (0–6.5)
Goldstuck (1983) ^b	UK	Cu 7, Cu 7 Mini, Cu T, MLCu 250, MLCu 250 short	10+	71	64	0	0 (0–5.6)
Guillebaud <i>et al.</i> (1983) ^b	UK	Cu 7, Lippes Loop	5	87	87	0	0 (0–4.2)
Gottardi <i>et al.</i> (1986) ^b	Italy	Cu T-200, Cu 7, Cu 7 Mini, MLCu-250	7	98	91	0	0 (0–4.0)
Luerti <i>et al.</i> (1986) ^b	Italy	MLCu-250, Cu 7, Cu T	7	117	102	0	0 (0–3.6)
Askalani <i>et al.</i> (1987) ^b	Egypt	Cu T-200	4	200	200	4	2.0 (0.5–5.0)
Yang <i>et al.</i> (1997)	China	ML Cu-375 SL	5	30	30	0	0 (0–11.6)
Li <i>et al.</i> (1999)	China	ML Cu-375 SL	5	101	101	0	0 (0–3.6)
Zhang <i>et al.</i> (1999)	China	ML Cu-375 SL	5	100	100	0	0 (0–3.6)
Lu (2000)	China	Copper IUD (unspecified)	5	28	28	0	0 (0–12.3)
Tian (2000)	China	Copper IUD (unspecified)	5	80	80	0	0 (0–4.5)
Li <i>et al.</i> (2001)	China	Cu T-380A	5	94	94	0	0 (0–3.8)
Zhao <i>et al.</i> (2001)	China	ML Cu-375 SL	5	98	98	1	1.0 (0.0–5.6)
Zhou and Xiao (2001) ^b	China	ML Cu-375 SL	5+	1013	998	1	0.1 (0.0–0.6)
Liu and Chen (2002a,b)	China	ML Cu-375 SL	5	80	80	1	1.2 (0.0–6.8)
Liu and Chen (2002a,b)	China	Copper IUD (unspecified)	3	95	95	0	0 (0–3.8)
Sun <i>et al.</i> (2003)	China	GyneFix	5	86	86	0	0 (0–4.2)
D'Souza <i>et al.</i> (2003) ^b	UK	GyneFix, Gyne-T 380S	Unspecified	173	169	0	0 (0–2.2)
Tang <i>et al.</i> (2003)	China	ML Cu-375 SL	5	98	98	1	1.0 (0.0–5.6)
Wang and Jiang (2003)	China	Copper IUD (unspecified)	5	86	86	0	0 (0–4.2)
Zhao and Wang (2004)	China	Cu T-220	5	108	108	0	0 (0–3.4)
Hou (2005)	China	Nova T-380	5	100	100	0	0 (0–3.6)
Li <i>et al.</i> (2005)	China	ML Cu-375 SL	5	150	150	0	0 (0–2.4)
Wen (2005)	China	Cu T-380A	5	218	218	1	0.5 (0.0–2.5)

Continued

Table I Continued

Study	Country	IUD type	Max. time to insertion (days)	Enrolled (n)	Efficacy population (n)	Pregnancies (n)	Pregnancy rate (%) (95% CI) ^a
Wang et al. (2006)	China	Cu T-380A	5	200	200	0	0 (0–1.8)
Dong and Wu (2007)	China	Copper IUD (unspecified)	5	87	87	0	0 (0–4.2)
Sheng and Zhang (2007)	China	Copper IUD (unspecified)	5	27	27	0	0 (0–12.8)
Song et al. (2007)	China	ML Cu-375 SL	5	180	180	0	0 (0–2.0)
Yang et al. (2008)	China	Cu T-380A	5	100	100	0	0 (0–3.6)
Hong (2008)	China	GyneFix	5	100	100	0	0 (0–3.6)
Ma et al. (2008)	China	MCu	5	26	26	0	0 (0–13.2)
Yang et al. (2008)	China	ML Cu-375 SL	5	86	86	0	0 (0–4.2)
He (2009)	China	Copper IUD (unspecified), ML Cu-375 SL	5	108	108	0	0 (0–3.3)
Hong (2009)	China	Copper IUD (unspecified)	5	70	70	0	0 (0–5.1)
Liu et al. (2010)	China	ML Cu-375 SL	5	162	162	1	0.6 (0.0–3.4)
Turok et al. (2010) ^b	USA	Cu T-380A	5	23	22	0	0 (0–15.4)
Wang (2010)	China	Copper IUD (unspecified)	5	40	40	0	0 (0–8.8)
Wu et al. (2010) ^b	China	Cu T-380A	5	1963	1893	0	0 (0–0.2)

^aWhere the pregnancy rate is zero, a one-sided 97.5% CI is calculated. Otherwise, a 95% exact binomial CI is calculated.

^bIndicates publication in English; otherwise, studies were published in Chinese.

days of unprotected intercourse, although one study included 18 (out of 998) insertions beyond 5 days (Zhou and Xiao, 2001), three studies provided insertions up to 7 days after intercourse (Gottardi et al., 1979; Lippes et al., 1979; Luerti et al., 1986), one allowed insertions up to 10 days (Black et al., 1980) and one included 24 insertions (out of 64) at 10 or more days post-coitus (Goldstuck, 1983). One study did not specify the time to insertion at all (D'Souza et al., 2003). The studies did not include sufficient information on the delay between intercourse and insertion of the IUD to enable us to analyze the data by delay.

Among 7034 post-coital IUD insertions, there were 10 pregnancies, for an overall failure rate of 0.14% (95% CI = 0.08–0.25%) (Table II). Six pregnancies occurred among 5629 subjects in the studies conducted in China (failure rate = 0.11%; 95% CI = 0.05–0.23%) and the remaining four pregnancies occurred among 200 subjects in one study conducted in Egypt (Askalani et al., 1987). Strikingly, this study is the only RCT with a non-treatment arm for a contraceptive product that we are aware of. Three hundred women who had engaged in unprotected intercourse around the time of ovulation (and so had a relatively high probability of pregnancy) were randomized to either post-coital insertion of a Cu T-200 or no treatment. The pregnancy rates were 2% among the treatment group and 22%

in the expectant management group. The failure rate in the treatment arm of this study is surprisingly high, and significantly higher than the rate in all other countries combined ($P = 0.0001$); in contrast, the results among the five countries excluding Egypt are homogeneous ($P = 1$). If the true failure rate in Egypt were the same as in the other five countries (0.000878), then the chance of observing four or more pregnancies is vanishingly small, ~ 1 in 30 000 ($P = 0.00004$). This high failure rate can possibly be explained by the fact that women were specifically selected if they had had intercourse around the time of ovulation; in any event Egypt is a clear outlier. If the unusual results from the Egypt study were excluded, the overall failure rate would be 0.09% (95% CI = 0.04–0.19%); this is our preferred estimate.

Discussion

Our data suggest that IUDs are a highly effective method of emergency contraception, with a failure rate of less than one per thousand. The copper IUD is by far the most effective emergency contraceptive option, followed by mid-dose mifepristone (25–50 mg) or ulipristal acetate (failure rate $\sim 1.4\%$) and then levonorgestrel (failure rate $\sim 2\text{--}3\%$) (Cheng et al., 2008; Glasier et al., 2010).

Table II Failure rates for IUD use as emergency contraception, by country.

Country	Population	Pregnancies	Rate (%)	Exact 95% CI (%)
China	5629	6	0.11	0.05 0.23
UK	496	0	0.00	0.00 0.70
USA	401	0	0.00	0.00 0.85
Italy	253	0	0.00	0.00 1.38
Egypt	200	4	2.00	0.69 5.03
The Netherlands	55	0	0.00	0.00 5.93
Total	7034	10	0.14	0.08 0.25
Total excluding Egypt	6834	6	0.09	0.04 0.19

Data from two randomized trials of the ulipristal acetate and levonorgestrel regimens suggest that the efficacy of levonorgestrel declines sharply as BMI increases. Statistical models indicate that, among women with a BMI of 26 kg/m² or higher presenting after unprotected intercourse, levonorgestrel is no more effective than no treatment. Ulipristal acetate appears to retain its efficacy at higher BMI levels but is no more effective than no treatment at a BMI of 35 or higher (Glasier *et al.*, 2011). There is no clinical concern about the loss of effectiveness of the IUD with an increase in BMI; therefore, an IUD would be a particularly good choice for obese women presenting after unprotected sex.

Several recent studies exploring the awareness of and the interest in IUDs among women seeking emergency contraception identified barriers to a greater use of IUDs including cost, waiting time (patients are not always able to get an IUD the day that they present for emergency contraception), low levels of awareness and understanding among patients and lack of participation among providers (Schwarz *et al.*, 2009; Bharadwaj *et al.*, 2011; Turok *et al.*, 2011; Wright *et al.*, 2012). Two of these studies found that around 12% of women presenting for emergency contraception or walk-in pregnancy testing would consider an IUD as an alternative to emergency contraceptive pills (Schwarz *et al.*, 2009; Turok *et al.*, 2011); this finding indicates the considerable potential to increase IUD uptake among women who have recently had unprotected sex. A study of contraceptive providers in CA, USA, showed that 85% of clinicians never recommended the IUD for emergency contraception, and 93% require at least two visits for an IUD insertion (Harper *et al.*, 2012).

Our review has several limitations. The initial intention of this study was to assess the efficacy of IUDs for emergency contraception by the day of insertion (how many days had elapsed between unprotected intercourse and insertion of the IUD), but the studies generally did not include sufficient detail about the day of insertion among the efficacy-evaluable population. Therefore, our analysis groups all cases together, regardless of the length of delay between intercourse and insertion of the IUD. Similarly, we are unable to provide estimates of the efficacy by parity, individual pregnancy risk (the cycle day on which intercourse occurred) and IUD type, as most studies did not provide detailed information on these variables. Finally, as in any review, it is possible that studies may have been unintentionally excluded owing to incomplete search results. We were not able to include publications in languages other than English and Chinese; it

is a limitation that relevant studies in other languages were omitted as these might have added strength to these results. However, we believe that English and Chinese journals include the majority of publications on this topic.

Conclusions

Despite the limitations of the original data, this study contributes to the literature and to clinical practice by providing the most comprehensive review to date of the efficacy of IUDs used for emergency contraception. These results provide clear evidence that IUDs are a highly effective method of emergency contraception, as 99.86% of users overall did not become pregnant after unprotected intercourse when an IUD was inserted post-coitally. When we exclude the data from the Egyptian study (which does not represent the typical clinical scenario because the investigators intentionally selected women who were at a greater risk of pregnancy), the failure rate is 0.09%.

The cost, clinical protocols and lack of awareness among both patients and providers are barriers to a greater uptake of IUDs for emergency contraception. Increasing the use of IUDs for emergency contraception is an important strategy for reducing an individual woman's chance of becoming pregnant after unprotected intercourse. In addition, if left in place for ongoing contraception, copper IUDs provide highly effective contraception for at least 10 years, and can contribute to decreasing unintended pregnancy rates over the long term. Therefore, we conclude that IUDs should be routinely included as an emergency contraceptive option whenever clinically feasible and appropriate.

Authors' roles

K.C. conducted the English literature search, abstracted data from the English literature and drafted the article. H.Z. conducted the Chinese literature search, abstracted data from the Chinese literature and provided comments on the manuscript. N.G. abstracted data from the English literature and provided comments on the manuscript. L.C. abstracted data from the Chinese literature and provided comments on the manuscript. J.T. oversaw the quantitative analysis and provided comments on the manuscript.

Funding

Support for Dr Trussell's work was provided by the Eunice Kennedy Shriver National Institute of Child Health and Development, Grant 2R24HD047879. No other external support was provided for this study.

Conflict of interest

None declared.

References

- ACOG Practice Bulletin. Clinical management guidelines for obstetrician-gynecologists. Number 112, emergency contraception. *Obstet Gynecol* 2010;**115**:1100–1108.
- Askalani AH, Al-Seniy AM, Al-Agizy HM, Salem HI, Al-Masry GI, El-Sadek SM. Evaluation of copper T-200 as a post-coital contraceptive. *Egypt Obstet Gynecol* 1987;**13**:63–66.
- Bharadwaj P, Saxton J, Mann S, Jungmann E, Stephenson J. What influences young women to choose between the emergency contraceptive pill and an intrauterine device? A qualitative study. *Eur J Contracept Reprod Health Care* 2011;**16**:201–209.
- Black TRL, Goldstuck ND, Spence A. Post-coital intrauterine device insertion—a further evaluation. *Contraception* 1980;**22**:653–658.
- Bromwich PD, Parsons AD. The establishment of a postcoital contraception service. *Br J Fam Plan* 1982;**8**:16–19.
- Centers for Disease Control and Prevention. U.S. Medical Eligibility Criteria for Contraceptive Use, 2010, **Vol. 59**. MMWR, 2010, 1–85. http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5904a1.htm?s_cid=rr5904a1_e.
- Cheng L, Guelmenzoglu AM, Piaggio G, Ezcurra E, Van LPFA. Interventions for emergency contraception. *Cochrane Database Syst Rev* 2008;**2**.
- Dean G, Schwarz EB. Intrauterine contraceptives. In: Hatcher RA, Trussell J, Nelson A, Cates W, Stewart F, Kowal D, Policar M (eds). *Contraceptive Technology*. New York, NY: Ardent Media, 2011.
- Dong SZ, Wu SH. Mifepristone, levonorgestrel and IUD for emergency contraception: a 268-case control study. *J Chin Mod Gyn Ob* 2007;**4**:127–129 [Chinese].
- D'Souza R, Masters T, Bounds W, Guillebaud J. Randomised controlled trial assessing the acceptability of GyneFix versus Gyne-T380S for emergency contraception. *J Fam Plan Reprod H* 2003;**29**:23–29.
- Faculty of Sexual & Reproductive Healthcare, Clinical Effectiveness Unit. Clinical Guidance on Emergency Contraception 2011. <http://www.fsrh.org/pdfs/CEUguidanceEmergencyContraception11.pdf>.
- Fan HM, Zhou LY. Emergency contraception with Multiload Cu-375SL IUD: a multicentre clinical trial. *J Reprod Med* 2001;**10**:70–77 [Chinese].
- Finer LB, Zolna MR. Unintended pregnancy in the United States: incidence and disparities 2006. *Contraception* 2011;**84**:478–485.
- Friedman EHI, Rowley DEM. Post-coital contraception—a two-year evaluation of a service. *Br J Fam Plan* 1987;**13**:139–144.
- Glasier AF, Cameron S, Fine P, Logan S, Casale W, Van Horn J, Sogor L, Blithe D, Scherrer B, Mathe H et al. Ulipristal acetate versus levonorgestrel for emergency contraception: a randomised non-inferiority trial and meta-analysis. *Lancet* 2010;**375**:555–562.
- Glasier A, Cameron ST, Blithe D, Scherrer B, Mathe H, Levy D, Gainer E, Ulmann A. Can we identify women at risk of pregnancy despite using emergency contraception? Data from randomized trials of ulipristal acetate and levonorgestrel. *Contraception* 2011;**84**:363–367.
- Goldstuck ND. Delayed postcoital IUD insertion. *Contracept Deliv Syst* 1983;**4**:293–296.
- Gottardi G, Marzi M, Pozzi S. Postcoital estrogen or IUD? *IPPF Euro Reg Bull* 1979;**8**:7–8.
- Gottardi G, Spreafico A, de Orchi L. The postcoital IUD as an effective continuing contraceptive method. *Contraception* 1986;**34**:549–558.
- Guillebaud J, Kubba A, Rowlands S, White J, Elder MG. Post-coital contraception with danazol, compared with an ethinyloestradiol-norgestrel combination or insertion of an intra-uterine device. *J Obstet Gynaecol* 1983;**3**(Suppl. 2):S64–S68.
- Harper CC, Speidel JJ, Drey EA, Trussell J, Blum M, Darney PD. Copper intrauterine device for emergency contraception: clinical practice among contraceptive providers. *Obstet Gynecol* 2012;**119**:220–226.
- He YM. Comparison of mifepristone and copper IUD for emergency contraception. *Shanxi Med J* 2009;**38**:703 [Chinese].
- Hong SZ. GyneFix for emergency contraception. *Contemp Med* 2008;**145**:36–37 [Chinese].
- Hong CP. A clinical study of emergency contraception methods: levonorgestrel and IUD. *Chin Health Care Nutr* 2009;**18**:91–92 [Chinese].
- Hou XL. Clinical analysis of NovaT-380 IUD as emergency contraception. *Chin Gen Prac* 2005;**8**:292–293 [Chinese].
- Hubacher D, Finer LB, Espey E. Renewed interest in intrauterine contraception in the United States: evidence and explanation. *Contraception* 2011;**83**:291–294.
- Hutchinson F. Experience at London brook advisory centres. In: Grahame H (ed). *Postcoital Contraception: Methods, Services and Prospects*. London: Pregnancy Advisory Service, 1983, 40–42.
- Lader D. Contraception and Sexual Health, 2008/09. *Opinions Survey Report* 2009;**41**.
- Li HZ, Wang P, Li SZ, Wu BS, Zuo WL. Clinical trial of the efficacy of the copper IUD for emergency contraception. *Chin J Imp Birth Outcomes Child Dev* 1999;**10**:157–159 [Chinese].
- Li H, Tong Q, Huang XJ, Shu LZ. TCu-380A IUD for emergency contraception (with a one-year follow-up available). *J Tibet Med* 2001;**22**:47–49 [Chinese].
- Li FX, Qian XJ, Wu WP. A case control study of active IUD and mifepristone for emergency contraception. *Prac Med J* 2005;**21**:2313–2314 [Chinese].
- Lippes J, Malik T, Tatum HJ. The postcoital copper-T. *Adv Plann Parent* 1976;**11**:24–29.
- Lippes J, Tatum H, Maulik D, Zielezny M. Postcoital copper IUDs. *Adv Plann Parent* 1979;**14**:87–94.
- Liu XJ. Multiload for emergency contraception. *Chin Prac Med* 2010;**30**:135–136 [Chinese].
- Liu LP, Chen AX. Comparison of emergency contraception with IUD and mifepristone. *J Changzhi Med Col* 2002a;**16**:198–199 [Chinese].
- Liu Y, Chen XX. IUD insertion and mifepristone for emergency contraception. *J Qiqihar Med Col* 2002b;**23**:890–891 [Chinese].
- Lu J. Clinical application of the copper IUD for emergency contraception. *Chin J Mod Med* 2000;**10**:48 [Chinese].
- Luerti M, Tonta A, Ferla P, Molla R, Santini F. Post-coital contraception by estrogen progestagen combination or IUD insertion. *Contraception* 1986;**33**:61–68.
- Ma XY, Zhang ZH, Sun CY. A report on 26 cases of MLCu IUD as emergency contraception. *Mod Prev Med* 2008;**35**:4372–4373 [Chinese].
- Moreau C, Lydie N, Warzwaski J, Bajos N. Activite sexuelle, infections sexuellement transmissibles, contraception. In: Beck F, Guilbert P, Gautier A (eds). *Barometre Sante* 2005. Saint Denis: IPNES, 2008, 329–353.
- Mosher WD, Jones J. Use of contraception in the United States: 1982–2008. *Vital Health Stat* 2010;**29**:1–44.

- 685 National Population and Family Planning Commission, Population and
Development Research Center of China. Nationwide distribution of
contraceptive methods in married reproductive age women. *Pop Fam
Plan Data* 2010. China Population Publishing House [Chinese]. p. 194.
- Salem RM. New attention to the IUD: expanding women's contraceptive
options to meet their needs. *Popul Rep B* 2006;**7**::1–26.
- 690 Schwarz EB, Kavanaugh M, Douglas E, Dubowitz T, Creinin MD. Interest in
intrauterine contraception among seekers of emergency contraception
and pregnancy testing. *Obstet Gynecol* 2009;**113**:833–839.
- Sheng QY, Zhang XL. A clinical study to compare mifepristone and IUD
for emergency contraception. *Chin Comm Doc* 2007;**9**:38 [Chinese].
- 695 Singh S, Wulf D, Hussain R, Bankole A, Sedgh G. Abortion Worldwide: A
Decade of Uneven Progress, 2009. [http://www.guttmacher.org/pubs/
Abortion-Worldwide.pdf](http://www.guttmacher.org/pubs/Abortion-Worldwide.pdf).
- Song CH, Wang Y, Chen P, Wang D, Lu WH. A case-control study of
mifepristone and IUD for emergency contraception. *Chin Med Factory
Mine* 2007;**20**:630–631 [Chinese].
- 700 Sun Q, Dong J, Sun Y. GyneFix IUD for emergency contraception.
J Shandong Med 2003;**43**:65 [Chinese].
- Tang LR, Nan XP, Wu MH, Fan HM. A clinical study of Multiload 375-SL for
emergency contraception. *Beijing Med J* 2003;**25**:137–138 [Chinese].
- Tian QH. Clinical trial of mifepristone and IUD for emergency
contraception. *J Henan Med Col for Staff and Workers* 2000;**12**:51
705 [Chinese].
- Trussell J, Lalla AM, Doan QV, Reyes E, Pinto L, Gricar J. Cost
effectiveness of contraceptives in the United States. *Contraception*
2009;**79**:5–14.
- 710 Turok DK, Gurtcheff SE, Handley E, Simonsen SE, Sok C, Murphy P. A
pilot study of the Copper T380A IUD and oral levonorgestrel for
emergency contraception. *Contraception* 2010;**82**:520–525.
- Turok DK, Gurtcheff SE, Handley E, Simonsen SE, Sok C. A survey of
women obtaining emergency contraception: are they interested in
using the copper IUD? *Contraception* 2011;**83**:441–446.
- 715 Tyrer LB. The copper-7 and postcoital contraception. *Adv Plann Parent*
1980;**15**:111–117.
- Van Santen MR, Haspels AA. Interception by post-coital IUD insertion.
Contracept Deliv Syst 1981;**2**:189–200.
- Wang XH. Copper IUD for emergency contraception: a case control
study. *Health Must-Read Mag* 2010;**6**:144 [Chinese].
- 720
- 725
- 730
- 735
- 740
- Wang JH, Jiang L. An 86-case analysis of copper IUD for emergency
contraception. *Ningxia Med J* 2003;**25**:353 [Chinese].
- Wang CP, Liu Y, Chang YF, Shao WQ. Feasibility of IUD and low-dose
mifepristone for emergency contraception. *J Fertility* 2006;**15**:271–273
745 [Chinese].
- Wen CL. Study of the efficacy of TCu380A for emergency contraception.
Prac Diag Treat J 2005;**19**:731–732 [Chinese].
- World Health Organization. *Medical Eligibility Criteria for Contraceptive Use*,
2009. [http://whqlibdoc.who.int/publications/2010/9789241563888_
eng.pdf](http://whqlibdoc.who.int/publications/2010/9789241563888_eng.pdf). 750
- Wright DW, Thompson PM. Monitoring a post-coital contraception
service. *Br J Fam Plan* 1986;**12**:88–91.
- Wright R, Frost C, Turok DK. A qualitative exploration of emergency
contraception users' willingness to select the copper IUD.
Contraception 2012;**85**:32–35. 755
- Wu S, Godfrey EM, Wojdyla D, Dong J, Cong J, Wang C, von Hertzen H.
Copper T380A intrauterine device for emergency contraception: a
prospective, multicentre, cohort clinical trial. *BJOG-Int J Obstet
Gynaecol* 2010;**117**:1205–1210.
- Yang MH. A clinical study of effect of the multiload as emergency
contraception. *Guide Chin Med* 2008;**6**:20–22 [Chinese]. 760
- Yang Y, Zhang XM, Wei DY, Yi SJ. Clinical study of mifepristone,
A-nordrin and intrauterine device used for emergency contraception.
J Reprod Med 1997;**6**:171–173 [Chinese].
- Yang D, Lei ZW, Li M, He Z, Xu LL. Clinical evaluation of emergency
contraception with TCu380A IUD. *Sichuan Med J* 2008;**29**:1–2
765 [Chinese].
- Zhang LP, Huang Q. Women's acceptance of IUDs as emergency
contraception. *Coll J Wannan Med Sch* 2005;**16**:117–118 [Chinese].
- Zhang LJ, Jing XP, Wen LJ. A case-control study of mifepristone and the
copper IUD for emergency contraception. *Chin J Obstet Gynecol* 1999;
770 **34**:569–570 [Chinese].
- Zhao YH, Wang XL. Copper IUD for emergency contraception. *Chin J
Med Res Appl* 2004;**2**:31–32 [Chinese].
- Zhao H, Tang LR, Wu MH, Cheng H. Clinical research on mifepristone
and IUD for emergency contraception. *J Cap Univ Med Sci* 2001;
775 **22**:273–274 [Chinese].
- Zhou L, Xiao B. Emergency contraception with multiload Cu-375SL IUD: a
multicenter clinical trial. *Contraception* 2001;**64**:107–112.
- 780
- 785
- 790
- 795